

On plucking the plant, the two leaves are found connected below, with an obscure flower in their fork, which had rested on the surface of the ground, or been buried under the decayed leaves and soil. Its flowering time is from May to July.

This plant, from the number of its stamens, is placed by Linnæus and Michaux in the class *Dodecandria*. Pursh, who has omitted this class in his Flora, has transferred the Asarums to *Gynandria*, from the circumstance that the stamens are inserted on the germ. This place however is not better suited to the Asarum, than to a multitude of other plants whose germ is inferior.

Linnæus' natural order for this plant is *Sarmentaceæ* and Jussieu's *Aristolochiæ*.

Generic character. *Calyx three or four cleft, superior; corolla none; anthers growing to the middle of the filaments. Capsule coriaceous, crowned.*

Specific character. *Leaves two, reniform; calyx woolly, cleft to the base; its segments spreading at top.*

The root of the Asarum is creeping, fleshy, and somewhat jointed. Leaves kidney shaped, pubescent on both sides, with long, round, hairy petioles. Flower solitary, growing from the fork of the stem, on a pendulous hairy peduncle. Ca-

lyx very hairy or woolly, consisting of three broad, concave leaflets, which are mostly of a brownish or dull purple on the inside at top and bottom, and terminated by a long, spreading, inflected point, with reflexed sides. The colour varies greatly according to the amount of light which the plant enjoys, being sometimes nearly green. Stamens twelve, inserted on the germ at a distance from the calyx, the alternate ones longer. Anthers growing to the filaments below their extremity. Near the divisions of the calyx are three short, curved, filamentary substances, which may perhaps be called nectaries. Germ inferior, somewhat hexagonal, marked at top inside with a dark red line; style conical, striate, parted at top into six recurved, radiating stigmas.

The root of the Asarum has an agreeable aromatic taste, which is intermediate between that of ginger and the aristolochia serpentaria. This quality has given it the names of *Wild ginger* and *Snake root* in different sections of the country. The name *Colt's foot* is also applied to it.

The chemical trials, to which I have subjected the root, bring to view the following substances:—

1. A light coloured, pungent, volatile oil, possessing the characteristic taste and smell of the plant in a high degree. 2. A resin, which is of a red-